

October 1996  
hep-th/9610252

IASSNS-HEP 96/108  
UPR-722-T

# Resolution of Cosmological Singularities

**Finn Larsen**<sup>1</sup>

*Department of Physics and Astronomy  
University of Pennsylvania  
Philadelphia, PA 19104*

and

**Frank Wilczek**<sup>2</sup>

*School of Natural Sciences  
Institute for Advanced Study  
Princeton, NJ 08540*

## Abstract

We show that a class of 3+1 dimensional Friedmann-Robertson-Walker cosmologies can be embedded within a variety of solutions of string theory. In some realizations the apparent singularities associated with the big bang or big crunch are resolved at non-singular horizons of higher-dimensional quasi-black hole solutions (with compactified *real* time); in others plausibly they are resolved at D-brane bound states having no conventional space-time interpretation.

---

<sup>1</sup>e-mail: larsen@cvtic.hep.upenn.edu

<sup>2</sup>e-mail: wilczek@sns.ias.edu

In the course of their historic treatment of the collapse of a cloud of pressureless dust in general relativity, Oppenheimer and Snyder introduced a striking construction: the joining of an external Schwarzschild solution to the collapse phase of a closed Friedmann-Robertson-Walker, homogeneous and isotropic model universe [1]. The possibility to connect a static exterior geometry with a dynamic interior occurs because the character – space-like *versus* time-like – of the Schwarzschild  $r$  and  $t$  variables changes upon passage through the horizon. Indeed in this way even the interior of a vacuum Schwarzschild black hole can be interpreted as a cosmological solution, although a highly anisotropic one. To be specific, if we define  $v \equiv \sqrt{8m}(2m - r)^{\frac{1}{2}}$  then slightly inside the horizon the line element takes the form

$$ds^2 = -dv^2 + \frac{v^2}{16m^2}dt^2 + (2m)^2 d\Omega_2^2, \quad (1)$$

which appears to represent a primitive crunch singularity of the spatial dimension  $t$  as  $v \rightarrow 0^+$ . In reality of course there is no curvature singularity, but only a singularity of the  $v$  ‘time’ coordinate that is resolved in the complete embedding.

Here we will discuss some analogous phenomena in string theory and simple effective field theories derived from it. As we shall see, in this framework several new features appear: the larger number of available space dimensions make it possible to embed a homogeneous and isotropic model 3+1 dimensional model universe in an interesting, non-singular fashion; the dilaton, moduli, and antisymmetric tensor fields can provide interesting forms of ‘matter’; and – perhaps most challenging – the possibility arises that cosmological singularities are resolved in an intrinsically string-theoretic way that has no conventional space-time interpretation as it involves D-branes and their open string dressing.

It is convenient to begin the discussion by reference to the five dimensional Schwarzschild metric

$$ds^2 = -(1 - \frac{\mu}{r^2})dt^2 + (1 - \frac{\mu}{r^2})^{-1}dr^2 + r^2 d\Omega_3^2 \quad (2)$$

Inside the horizon it can be written after obvious renaming of variables as

$$ds^2 = e^{-4\sigma} dy^2 - e^{4\sigma} d\tau^2 + \tau^2 d\Omega_3^2 \quad (3)$$

$$e^{-4\sigma} = \frac{\mu}{\tau^2} - 1 \quad (4)$$

The range of  $\tau$  is  $0 < \tau^2 < \mu$ . To obtain an interesting embedded cosmology along the lines mentioned above, we compactify  $y$ . (Such a construction was also employed in [2, 3].) This induces an effective 4-dimensional dilaton  $e^{-2\phi_4} = e^{-2\sigma}$  that can be interpreted as an independent matter field. In string frame the dimensionally reduced metric is eq. 3, with the  $dy^2$  term omitted. In terms of the Einstein frame metric  $g_{E\mu\nu} = e^{-2\phi_4} g_{S\mu\nu}$  the four dimensional cosmology becomes

$$ds_E^2 = -e^{2\sigma} d\tau^2 + e^{-2\sigma} \tau^2 d\Omega_3^2 \quad (5)$$

with  $e^{-2\sigma}$  as in eq. 4. We regard the Einstein frame line element as privileged, both because it governs the response of conventional test bodies and because it is invariant under the standard duality transformations. Friedmann coordinates are introduced as

$$ds_E^2 = -dt^2 + R(t)^2 d\Omega_3^2 \quad (6)$$

The comoving time  $t$  spans a finite range and the scale factor  $R(t) \simeq |t - t_{i,f}|^{\frac{1}{3}}$  at early and late times. Kaluza-Klein compactification of Einstein gravity in  $D = 5$  gives the effective  $D = 4$  Lagrangian. It is a free scalar, minimally coupled to gravity:

$$16\pi G_N \mathcal{L} = R_E^{(4)} - 6g^{\mu\nu} \partial_\mu \sigma \partial_\nu \sigma \quad (7)$$

For a homogeneous field the ensuing energy momentum tensor describes matter with  $\rho = p$ . Friedmann's equations with this equation of state of course reproduce the  $|t - t_{i,f}|^{\frac{1}{3}}$  behavior found at early and late times.

Thus the  $D = 5$  vacuum black hole solution with a coordinate singularity – the horizon – and a physical singularity – the origin – becomes  $D = 4$  cosmology with scalar matter and physical (curvature) singularities at both initial and final times. (For a related discussion of certain extremal black hole singularities see [4].) From the perspective of the interior cosmology, either direction of temporal flow is equally good. The compactification of  $y$  appears innocuous for the interior solution, but its continuation beyond the horizon involves compactifying the real time  $t$  ( $=y$ ), so that there are closed time-like paths and normal causality conditions are violated in the exterior. Although these violations should not be accepted lightly, note that they are restricted to the necessarily exotic epoch ‘after’ the end of time. In any case, since they

arise in the natural continuation of an apparently reasonable cosmology, to forbid them would involve a new cosmological veto. At the horizon,  $t$  is not a good coordinate and one must analyze using (say) Kruskal coordinates  $U, V$ . The periodic identification of  $t$  becomes a re-scaling  $U \rightarrow \alpha U, V \rightarrow \frac{1}{\alpha} V$ . Though there is no curvature singularity, there is an unusual non-Hausdorff topology at the fixed point  $U = V = 0$ .

A recurrent theme in recent work on duality is that the low energy field theory develops singularities when heavy states of the full theory become light [5, 6, 7]. In the re-interpretation of big crunch as a higher-dimensional horizon the relevant heavy states are the Kaluza-Klein modes from the compactified dimension: at late times they become light and important for the dynamics; ultimately, they take over the role of time.

This simplest model has several deficiencies, including a remaining singularity at the origin of the  $D = 5$  black hole. In the following we generalize the model and

- embed the cosmology in a solution that recognizably derives from string theory.
- determine whether cosmologies derived from string theory naturally involve other kinds of matter.
- exhibit solutions in which the singularities are resolved by a new scenario, using that D-branes are singular as classical solutions but non-singular in their physical behavior. Here the apparent singularity of the low energy field theory arises because open string excitations, ignored in the effective field theory, become light close to the D-branes.

Large classes of cosmological solutions, including some of those considered below - but not their embedding inside black holes - have been analyzed recently [8, 9, 10, 11]. Configurations with non-trivial RR-fields are of particular interest because they can be represented as simple conformal field theories [12]. The starting point is the classical solution associated with the extremal Dirichlet p-brane [13] (for a review of string solitons see [14]):

$$ds_S^2 = e^{2\xi}(-dy^2 + dx_1^2 + \cdots + dx_p^2) + e^{-2\xi}(dx_{p+1}^2 + \cdots + dx_9^2) \quad (8)$$

$$e^{-2\phi_{10}} = e^{-2\xi(p-3)} \quad (9)$$

$$F_{p+2} = \partial_\mu e^{4\xi} dy \wedge dx_1 \wedge \cdots \wedge dx_p \wedge dx^\mu \quad (10)$$

$$e^{-4\xi} = 1 + \frac{q}{r^2}, \quad (11)$$

where  $F_{p+2}$  is the  $p+2$  form RR-field strength. We have written the solution in the  $D=10$  string frame, and anticipating compactification to  $D=5$  have arranged a periodic array in 5 small spatial dimensions. Then the harmonic function  $e^{-4\xi}$  assumes the form  $r^{-2}$  where  $r$  is a spatial 4-vector for any  $p$  ( $\leq 5$ ). The extremal metric eq. 8 has no horizon and therefore no interior; so we need its ‘black’ generalization [15, 16]. Inside the horizon it reads after renaming of variables

$$ds_S^2 = e^{2\xi}(e^{-4\sigma} dy^2 + dx_1^2 + \cdots + dx_p^2) + e^{-2\xi}(dx_{p+1}^2 + \cdots + dx_5^2) + e^{-2\xi}(-e^{4\sigma} d\tau^2 + \tau^2 d\Omega_3^2), \quad (12)$$

where  $e^{-4\xi} = \frac{q}{\tau^2} + 1$  and as before  $e^{-4\sigma} = \frac{\mu}{\tau^2} - 1$ , with  $\mu$  a numerical ‘non-extremality’ parameter. The field strength is also modified, as we will discuss presently. Compactification – including compactification of  $y$  – introduces an effective dilaton  $e^{-2\phi_4} = e^{2\xi-2\sigma}$  for all  $p$ . The  $D=4$  Einstein metric again becomes eq. 5. Thus the  $D$ -brane generates the same interior space-time as the Schwarzschild solution!

To understand this result in more detail, consider the metric eq. 12 an *ansatz* with the  $D=4$  fields  $\sigma$  and  $\xi$  undetermined *a priori*. Similarly, use eq. 9 as an *ansatz* for the dilaton and introduce an independent scalar field  $A$  that arises because the  $(p+2)$ -form field strength reduces to a 1-form, written as  $\nabla A$ . Then the  $D=10$  string action

$$16\pi G_N \mathcal{L} = e^{-2\phi_{10}} [R_S^{(10)} + 4(\nabla\phi_{10})^2] - \frac{1}{2(p+2)!} F_{p+2}^2 \quad (13)$$

gives rise to the effective  $D=4$  Einstein frame action

$$16\pi G_N \mathcal{L} = R_E^{(4)} - 6(\nabla\sigma)^2 - 8(\nabla\xi)^2 + 8\nabla\xi\nabla\sigma - \frac{1}{2}e^{4\sigma-8\xi}(\nabla A)^2 \quad (14)$$

after some calculation. The stress-energy tensor corresponding to this Lagrangian describes  $\rho = p$  matter, just as was the case without  $D$ -branes. The equations of motion for the scalar fields can be written

$$\nabla^2\sigma = 0 \quad (15)$$

$$\nabla_\mu(e^{4\sigma-8\xi}\nabla^\mu A) = 0 \quad (16)$$

$$4\nabla^2\xi + e^{4\sigma-8\xi}(\nabla A)^2 = 0. \quad (17)$$

We expect the solution

$$ds_E^2 = -e^{2\sigma} d\tau^2 + e^{-2\sigma} \tau^2 d\Omega_3^2 \quad (18)$$

$$e^{-4\sigma} = \frac{\mu}{\tau^2} - 1 \quad (19)$$

$$\partial_\tau A = \frac{2Q}{\tau^3} e^{8\xi} \ ; \ Q^2 = q(q + \mu) \quad (20)$$

$$e^{-4\xi} = \frac{q}{\tau^2} + 1 \ . \quad (21)$$

It is easy to verify that eqs. 18-19 indeed satisfies eq. 15 and that eq. 16 requires  $\partial_\tau A \propto \partial_\tau e^{4\xi}$  – which is precisely the *ansatz* eq. 20 above, with the proportionality constant, the charge, parametrized by  $Q$ . The final equation eq. 17 is non-trivial and is satisfied by virtue of cancellations that appear rather mysterious. Indeed in the energy-momentum tensor

$$16\pi G_N T_{\tau\tau} = 6\dot{\sigma}^2 + 8\dot{\xi}^2 - 8\dot{\xi}\dot{\sigma} + \frac{1}{2}e^{4\sigma-8\xi}\dot{A}^2 \quad (22)$$

the first term is sufficient to satisfy the Friedmann equations because  $\xi = 0$  is the solution without the D-brane. Therefore the subsequent terms must – and do – cancel. The higher dimensional physics leads us to define the scalar fields in such a way that only one of them couples to gravity, at least on-shell and for homogeneous solutions. The additional fields are somewhat marginal from the  $D = 4$  perspective, in that they carry no net energy, but they parametrize inequivalent embeddings of the  $D = 4$  theory into a higher dimensional theory.

The  $D = 4$  effective action eq. 14 is independent of  $p$  but the relation between the scalar fields  $\xi$  and  $\sigma$  and the higher dimensional geometry does depend on  $p$ . It is straightforward to verify explicitly that branes with coupling to the NS fields are similarly described by the action eq. 14, with yet another higher dimensional interpretation of the scalar fields. In fact, U-duality of the type II string theory implies that *all* black branes must give the same  $D = 4$  Einstein metric.

In all the models discussed so far, the big crunch (or big bang) singularity is resolved as a horizon in  $D = 5$  but the big bang (or big crunch) singularity remains a curvature singularity. In the microscopic theory for extremal branes, the source is a very large number of coincident D-branes. The non-extremality parameter  $\mu$  can be introduced by considering non-supersymmetric excitations of this object [17, 18]. The physical nature of

the apparent singularity can be explored using test D-brane (or closed string) probes. Formally, the interactions between the probe and the background are governed by an annulus diagram. It is finite for  $\mu = 0$  due to supersymmetry.  $\mu \neq 0$  can be modelled by including appropriate vertex operators on the boundary of the annulus, and the diagram remains finite. Higher order corrections are also finite order by order, though not necessarily smaller. The situation seems analogous to the case where supersymmetry is broken by finite velocity [7]. In that case, at large distances it is legitimate to truncate the implied sum over string states and describe the interaction as the field theoretic one expressed in the classical solution, while at small distances and very small velocities the interaction can be described in terms of exchange of a few open string states. We expect that in our case  $\mu \neq 0$ , as for velocities that are not small, the interaction remains non-singular though at present it is not calculable.

Regular black holes in  $D = 5$  are obtained by combining three black branes [18, 19]. In fact, this solution is unique up to duality [20]. As a representative of this class, consider an NS string winding around the 1 direction, a  $D4$ -brane wrapped around the 2345 directions, and a  $D0$  brane. The line element in  $D = 10$  string frame form of these three intersecting black branes is

$$\begin{aligned} ds_S^2 &= e^{4\xi_w + 2\xi_0 + 2\xi_4 - 4\sigma} dy^2 + e^{4\xi_w - 2\xi_0 - 2\xi_4} dx_1^2 + \\ &+ e^{2\xi_4 - 2\xi_0} (dx_2^2 + dx_3^2 + dx_4^2 + dx_5^2) + e^{-2\xi_0 - 2\xi_4} (-e^{4\sigma} d\tau^2 + \tau^2 d\Omega_3^2) \\ e^{-2\phi_{10}} &= e^{6\xi_0 - 2\xi_4 - 4\xi_w} \end{aligned} \quad (23)$$

where as before  $e^{-4\sigma}$  and  $e^{-4\xi_i}$  are harmonic functions. (Here  $i = 1, 2, 3$ , by abuse of notation.) Alternatively, consider this an *ansatz* and derive the effective  $D = 4$  action

$$16\pi G_N \mathcal{L}_E^{(4)} = R_E^{(4)} - 6(\nabla\sigma)^2 + \sum_{i=1}^3 [8\nabla\sigma\nabla\xi_i - 8(\nabla\xi_i)^2 - \frac{1}{2}e^{4\sigma - 8\xi_i}(\nabla A_i)^2] \quad (24)$$

It is a consequence of U-duality that the fields  $\xi_i$  appear symmetrically despite the *ansatz* eq. 23. The equations of motion factorize appropriately so that their solution indeed becomes the expected one, *i.e.* eqs. 18–21 (with three identical copies of eqs. 20–21) .

To make contact with more familiar ideas, let us consider the special case

where the 3 charges are equal. Then upon reducing to 5 dimensions one finds

$$ds^2 = e^{8\xi-4\sigma} dy^2 + e^{-4\xi}(-e^{4\sigma}d\tau^2 + \tau^2 d\Omega_3^2) \quad (25)$$

and a time independent dilaton; so string frame agrees with Einstein frame. Introducing black hole notation the line element becomes

$$ds^2 = -\frac{1}{(1+\frac{q}{r^2})^2}(1-\frac{\mu}{r^2})dt^2 + (1+\frac{q}{r^2})\frac{1}{1-\frac{\mu}{r^2}}dr^2 + (1+\frac{q}{r^2})r^2 d\Omega_3^2, \quad (26)$$

and finally in terms of the geometrical radius  $R^2 = r^2 + q$  (which governs the volume of the overlying spheres)

$$ds^2 = -(1-\frac{R_+^2}{R^2})(1-\frac{R_-^2}{R^2})dt^2 + \frac{1}{(1-\frac{R_+^2}{R^2})(1-\frac{R_-^2}{R^2})}dR^2 + R^2 d\Omega_3^2, \quad (27)$$

where  $R_+^2 = q + \mu$  and  $R_-^2 = q$  specify the outer and inner horizons respectively. Also  $2G_N M = 2q + \mu$  and  $R_+^2 R_-^2 = Q^2$ . In this form it is clear that the horizons are indeed coordinate singularities and that the origin  $R = 0$  is a curvature singularity, even in the extremal case.

Returning to the general case the  $D = 4$  cosmological geometry is again the familiar one, which displays remarkable robustness. The dilaton behaves as  $e^{-2\phi_4} = e^{2\xi_0+2\xi_4-2\sigma}$  and approaches zero at the inner horizon/origin of the universe. This suggests that the coupling constant for string perturbation theory diverges. It appears now, however, to be a pseudo-singularity caused by an illegal truncation of the theory: specifically, the volume of the universe goes to zero because the length of the 5th dimension goes to zero. In  $D = 5$  all curvatures remain bounded and indeed small near the horizon provided  $\mu \gg l_P^2$  and  $q_i \gg l_P^2$  where  $l_P$  is the  $D = 5$  Planck length. The first condition is automatic for a macroscopic universe. The second requires that the inner horizon is well separated from the singularities. It will be satisfied when all the  $Q_i$  are large. If it is not satisfied, the singularity is not resolved within the field theory approach. However, it is plausible that the space-time singularity, which acts as the source of mass and charge, marks the approximate space-time locus (to the extent this notion makes sense) of the region where the brane intersection resides, and where an adequate description of the physics must take into account the open-string degrees of freedom which become light there.

The proposed cosmology is one where, at early times, the evolution is dominated by the Lagrangian eq. 24. Because of their stiff equation of state, the effective scalars will tend to dominate other forms of matter close to the apparent singularities. One or more of these scalars diverge close to the origin of time; so the  $D = 4$  effective field theory breaks down and must be replaced by something more exotic. Depending on the details (the number of non-vanishing  $\xi_i$ ) the natural description becomes  $D = 5$  effective field theory with closed time-like loops, or an open string effective field theory without a conventional interpretation in terms of geometric quantities. In both cases the transition is perfectly smooth.

Our models are quite schematic. Even if one of them were to give a valid first indication for physics very near the apparent big bang singularity, much more complex dynamics will come into play afterwards. As a simple example of the possibilities, note that if the crossover at late times is before the integration constants  $\pm 1$  in the harmonic functions become significant in the metric the universe could still be flat, or even open. The relevant solutions generalize those found in [2]:

$$ds_E^2 = -e^{2\sigma} d\tau^2 + e^{-2\sigma} \tau^2 d\Omega_3^2 \quad (28)$$

$$e^{-4\sigma} = \frac{\mu}{\tau^2} - k \quad (29)$$

$$e^{-4\xi_i} = \frac{q_i}{\tau^2} + 1 \quad (30)$$

$$\partial_\tau A_i = \frac{2Q_i}{\tau^3} e^{8\xi} \quad ; \quad Q_i^2 = q_i(kq_i + \mu) \quad (31)$$

Here  $k = -1, 0, 1$ . The solutions with  $k = -1, 0$  can not be interpreted as black holes.

**Acknowledgments:** We would like to thank A. Lukas and B. Ovrut for discussions. F.L. is supported in part by DOE grant DE-AC02-76-ERO-3071. F.W. is supported in part by DOE grant DE-FG02-90ER-40542

## References

- [1] J. R. Oppenheimer and H. Snyder. On continued gravitational contraction. *Phys. Rev. D*, 55:374–381, 1939.

- [2] K. Behrndt and S. Förste. String Kaluza-Klein cosmology. *Nucl. Phys. B*, 430:441–459, 1994. hep-th/9403179.
- [3] R. Poppe and S. Schwager. String Kaluza-Klein cosmologies with RR-fields. hep-th/9610166.
- [4] G. Gibbons, G. Horowitz, and P. Townsend. Higher-dimensional resolution of dilatonic black hole singularities. *Class. Quant. Grav.*, 12:297–318, 1995. hep-th/9410073.
- [5] N. Seiberg and E. Witten. Electric-magnetic duality in supersymmetric non-abelian gauge theories. *Nucl. Phys. B*, 426:19–52, 1994. hep-th/9407087.
- [6] B. R. Greene, D. Morrison, and A. Strominger. Black hole condensation and the unification of string vacua. *Nucl. Phys. B*, 451:109–120, 1995. hep-th/9504145.
- [7] M. Douglas, D. Kabat, P. Pouliot, and S. Shenker. D-branes and short distances in string theory. hep-th/9608024.
- [8] A. Lukas, B. Ovrut, and D. Waldram. Cosmological solutions of type II string theory. hep-th/9608195.
- [9] N. Kaloper. Stringy Toda Cosmologies. hep-th/9609087.
- [10] H. Lü, S. Mukherji, C. Pope, and K.-W. Xu. Cosmological solutions in string theories. hep-th/9610107.
- [11] A. Lukas, B. Ovrut, and D. Waldram. String and M-Theory Cosmological solutions with Ramond forms. hep-th/9610238.
- [12] J. Polchinski. Dirichlet-branes and Ramond-Ramond charges. *Phys. Rev. Lett.*, 75:4724, 1995. hep-th/9510017.
- [13] G. Horowitz and A. Strominger. Black strings and p-branes. *Nucl. Phys. B*, 360:197–209, 1991.
- [14] M. Duff, R. Khuri, and J. Lu. String Solitons. *Phys. Rept.*, 259:213–326, 1995. hep-th/9412184.

- [15] M. Duff, H. Lü, and C. Pope. The extremal branes of M-theory. *Phys. Lett. B*, 382:73–80, 1996. hep-th/9604052.
- [16] A. Tseytlin and M. Cvetič. Nonextreme black holes from nonextreme intersecting M-branes. hep-th/9606033.
- [17] C. Callan and J. Maldacena. D-brane approach to black hole quantum mechanics. hep-th/9602043, 1996.
- [18] G. Horowitz, J. Maldacena, and A. Strominger. Nonextremal black hole microstates and U-duality. *Phys. Lett. B*, 383:151–159, 1996. hep-th/9603109.
- [19] M. Cvetič and D. Youm. General rotating five-dimensional black holes of toroidally compactified heterotic string. *Nucl. Phys. B*, 476:118–132, 1996. hep-th/9603100.
- [20] C. Hull and M. Cvetič. Black holes and U-duality. hep-th/9606193.